

Forum Original Research Communication

The Role of Microglia in Paraquat-Induced Dopaminergic Neurotoxicity

XUE-FEI WU,^{1,2} MICHELLE L. BLOCK,¹ WEI ZHANG,^{1,3} LIYA QIN,¹ BELINDA WILSON,¹
WAN-QIN ZHANG,² BELLINA VERONESI,⁴ and JAU-SHYONG HONG¹

ABSTRACT

The herbicide paraquat (PQ) has been implicated as a potential risk factor for the development of Parkinson's disease. In this study, PQ (0.5–1 μ M) was shown to be selectively toxic to dopaminergic (DA) neurons through the activation of microglial NADPH oxidase and the generation of superoxide. Neuron-glia cultures exposed to PQ exhibited a decrease in DA uptake and a decline in the number of tyrosine hydroxylase-immunoreactive cells. The selectivity of PQ for DA neurons was confirmed when PQ failed to alter γ -aminobutyric acid uptake in neuron-glia cultures. Microglia-depleted cultures exposed to 1 μ M PQ failed to demonstrate a reduction in DA uptake, identifying microglia as the critical cell type mediating PQ neurotoxicity. Neuron-glia cultures treated with PQ failed to generate tumor necrosis factor- α and nitric oxide. However, microglia-enriched cultures exposed to PQ produced extracellular superoxide, supporting the notion that microglia are a source of PQ-derived oxidative stress. Neuron-glia cultures from NADPH oxidase-deficient (PHOX^{-/-}) mice, which lack the functional catalytic subunit of NADPH oxidase and are unable to produce the respiratory burst, failed to show neurotoxicity in response to PQ, in contrast to PHOX^{+/+} mice. Here we report a novel mechanism of PQ-induced oxidative stress, where at lower doses, the indirect insult generated from microglial NADPH oxidase is the essential factor mediating DA neurotoxicity. *Antioxid. Redox Signal.* 7, 654–661.

INTRODUCTION

PARKINSON'S DISEASE (PD) is a common neurodegenerative disorder characterized by the progressive and selective loss of dopaminergic (DA) neurons in the substantia nigra (SN) pars compacta (28), where >90% of PD cases are sporadic with unknown etiology (15). Epidemiological studies indicate that rural residence, farming, well water drinking, and exposure to agricultural chemicals are associated with an increased risk of developing PD (18, 31, 35, 36). In addition, exposure to several classes of environmental toxins, such as paraquat (PQ; 1,1'-dimethyl-4,4'-bipyri-

dinium), maneb, rotenone, and dieldrin, has resulted in DA neurotoxicity in animal models (20). Across this list of environmental neurotoxins, the common theme governing the selective DA neurotoxicity is the generation of oxidative stress. The following study sought to elucidate the source and the mechanism of PQ-induced oxidative stress at low concentrations (up to 1 μ M) and the consequent DA neurotoxicity.

Initially, the herbicide PQ was speculated to be toxic to DA neurons because of its structural similarity to 1-methyl-4-phenylpyridinium (MPP⁺), the active metabolite of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), the well known

¹Neuropharmacology Section, Laboratory of Pharmacology and Chemistry, National Institute of Environmental Health Sciences, Research Triangle Park, NC.

²Department of Physiology and ³Department of Neurology, First Clinical Hospital, Dalian Medical University, Dalian, China.

⁴Neurotoxicology Division, Office of Research and Development, National Health and Environmental Effects Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, NC.

parkinsonism-inducing agent discovered in synthetic heroin (16). Indeed, epidemiological studies have shown a strong correlation between the incidence of PD and exposure to PQ (18, 34). Studies using animal models have also supported the role of PQ and DA cell death, where intranigral injection of PQ induced dopamine (DA) depletion in the striatum, loss of DA neurons, glial reaction in the SN, and behavior abnormality (17). The effect of systemically administered PQ on the nigrostriatal system has been less consistent. Some studies have reported that PQ failed to damage DA neurons in SN and failed to deplete DA in striatum (29, 45, 46). However, more studies have indicated that systemic PQ exposure *in vivo* results in the selective degeneration of DA neurons in SN, DA depletion in striatum, and behavior abnormality (2, 6, 25, 26, 38, 39). Animals exposed to PQ combined with other pesticides also show selective DA neurotoxicity in the SN and behavioral deficits (1, 43). The abilities of PQ to induce oxidative stress in the brain (41, 47), to be taken up by DA neurons through the DA transporter (37, 38), and to up-regulate α -synuclein expression and facilitate its fibrillation (25, 44) may contribute to its DA neurotoxicity. However, the detailed mechanism governing PQ-induced DA neurotoxicity remains unclear.

There is increasing evidence supporting the association of inflammation and PD (20, 27), where microglia have been identified as the pivotal cell type governing inflammation-mediated neurodegeneration (20). Microglia are the resident immune cells of the central nervous system and contribute to neurodegeneration through the release of a variety of neurotoxic factors, including cytokines, free radicals, inflammatory prostaglandins, and fatty acid metabolites (17, 20, 24, 26, 42). Previous studies have observed a glial response in the midbrain of animals exposed to PQ (17, 26). However, the specific role of inflammation in PQ-induced DA neurotoxicity has yet to be investigated.

In this study, selective DA neurotoxicity was observed in rat and mouse mesencephalic neuron-glia cultures exposed to low (up to 1 μ M) concentrations of PQ. Here, we show that PQ-induced neurotoxicity was mediated through the presence of microglia and the activation of NADPH oxidase, which resulted in the production of the extracellular reactive oxygen species (ROS) responsible for DA neurotoxicity.

MATERIALS AND METHODS

Animals

All animals were treated in strict accordance with the National Institutes of Health (Bethesda, MD, U.S.A.) *Guide for Humane Care and Use of Laboratory Animals*. Timed pregnant (gestational day 14) adult female Fischer 344 rats were purchased from Charles River Laboratories (Raleigh, NC, U.S.A.). Eight-week-old male and female C57BL/6J (PHOX^{+/+}) and B6.129S6-Cybb^{tm1Din} (PHOX^{-/-}) mice were obtained from Jackson Laboratory (Bar Harbor, ME, U.S.A.) and maintained in a strict pathogen-free environment. Breeding of the mice was designed to achieve accurate timed pregnancy \pm 0.5 days. The PHOX^{-/-} mice lack the functional catalytic subunit of the NADPH oxidase complex, gp91. Be-

cause the PHOX^{-/-} mutation is maintained in the C57BL background, the C57BL/6J (PHOX^{+/+}) mice were used as control animals.

Reagents

Lipopolysaccharide (LPS; strain O111:B4) was purchased from Calbiochem (San Diego, CA, U.S.A.). Cell culture ingredients were obtained from Life Technologies (Grand Island, NY, U.S.A.). [³H]DA (28 Ci/mmol) and γ -[2,3-³H]aminobutyric acid (GABA; 81 Ci/mmol) were purchased from NEN Life Science (Boston, MA, U.S.A.). The polyclonal antibody against tyrosine hydroxylase (TH) was a kind gift from Dr. John Reinhard of Glaxo Smith Kline (Research Triangle Park, NC, U.S.A.). The monoclonal antibody raised against the CR3 complement receptor (OX42) was obtained from Pharmingen (San Diego, CA, U.S.A.). The biotinylated horse anti-mouse and goat anti-rabbit secondary antibodies were purchased from Vector Laboratories (Burlingame, CA, U.S.A.). WST-1 was purchased from Dojindo Laboratories (Gaithersburg, MD, U.S.A.). Tumor necrosis factor- α (TNF α) enzyme-linked immunosorbent assay (ELISA) kits were purchased from R&D Systems Inc. (Minneapolis, MN, U.S.A.). All other reagents came from Sigma-Aldrich Chemical Co. (St. Louis, MO, U.S.A.).

Mesencephalic neuron-glia cultures

Rat and mouse ventral mesencephalic neuron-glia cultures were prepared using a previously described protocol (22). In brief, mesencephalic tissues were dissected from embryonic day 14 Fischer 344 rats and PHOX^{+/+} or PHOX^{-/-} mice. Cells were dissociated via gentle mechanical trituration in minimum essential medium, immediately seeded (5×10^5 /well) in poly-D-lysine (20 μ g/ml) precoated 24-well plates, and maintained at 37°C in a humidified atmosphere of 5% CO₂ and 95% air. Cells were seeded in maintenance medium and treated with the treatment medium described previously (22). Three days after seeding, the cells were replenished with 0.5 ml of fresh maintenance medium. Cultures were treated 7 days after seeding.

Microglia-enriched cultures

Primary mixed microglia cultures were prepared from the whole brains of 1-day-old Fischer 344 rat pups, following the procedure described previously (23). Two weeks after seeding when the cells reached confluence, microglia were separated from astrocytes by shaking the flasks for 5 h at 150 rpm and replated at 1×10^5 in a 96-well plate. Enriched microglia were treated 24 h after seeding.

Mesencephalic microglia-depleted cultures

Microglia were depleted from mesencephalic neuron-glia cultures by the addition of 2 mM L-leucine methyl ester to the cultures 24 h after seeding. The cultures were exposed to L-leucine methyl ester for 72 h. Microglia-depleted cultures stained with OX-42 antibody showed <0.1% microglia.

Uptake assays (DA and GABA)

The DA and GABA uptake assays were performed as previously reported (8). In brief, after rinsing with warm Krebs–Ringer buffer, cultures were incubated for 20 min at 37°C with either 5 μ M [3 H]GABA or 1 μ M [3 H]DA. After incubation, cells were rinsed three times with 1 ml/well ice-cold Krebs–Ringer buffer, then lysed with 0.5 ml/well 1M NaOH, and mixed with 15 ml of scintillation fluid. Radioactivity was measured on a scintillation counter, where specific [3 H]GABA or [3 H]DA uptake was calculated by subtracting the mazindole or the NO-711 and β -alanine counts from the wells without the uptake inhibitors.

Immunostaining

Immunocytochemistry was performed using the conditions specified previously (21). DA neurons were identified by staining with the polyclonal antibody against TH. Microglia were stained with the monoclonal antibody raised against the CR3 receptor, OX-42. To quantify cell numbers, nine representative areas per well in the 24-well plate were counted under the microscope at 100 \times magnification by three individuals. The average of these scores was reported.

Superoxide assay

Extracellular superoxide ($O_2^{\cdot-}$) production from microglia was determined as reported previously (32) by measuring the superoxide dismutase (SOD)-inhibitable reduction of tetrazolium salt, WST-1 (19, 30, 40). In brief, immediately before treatment, enriched microglia were washed twice with Hanks' balanced salt solution (HBSS). To each well, 100 μ l of HBSS with or without SOD (600 U/ml), 50 μ l of vehicle, PQ, or LPS, and 50 μ l of WST-1 (1 mM) in HBSS were added. The cultures were incubated for 30 min at 37°C, and the absorbance at 450 nm was read with a SpectraMax Plus microplate spectrophotometer (Molecular Devices, Sunnyvale, CA, U.S.A.). The amount of SOD-inhibitable $O_2^{\cdot-}$ was calculated and expressed as percentage of vehicle-treated control cultures. Cell-free experiments determined that PQ treatment alone did not affect absorbance.

TNF α assay

The production of TNF α was measured with a commercial ELISA kit from R&D Systems, as described previously (21).

Nitrite assay

As an indicator of nitric oxide production, the amount of nitrite accumulated in culture supernatant was determined with a colorimetric assay using Griess reagent [1% sulfanilamide, 2.5% H_3PO_4 , 0.1% *N*-(1-naphthyl)ethylenediamine dihydrochloride] (11), as previously reported (21).

Statistical analysis

The data are expressed as the means \pm SEM, and statistical significance was assessed with an analysis of variance followed by Bonferroni's *t* test. A value of $p < 0.05$ was considered statistically significant.

RESULTS

PQ (0.5–1 μ M) is toxic to DA neurons

The DA neurotoxicity of PQ was measured in rat mesencephalic neuron-glia cultures. The ability of cells to take up [3 H]DA was measured to assess the effect of PQ on DA neuronal function. Exposure of neuron-glia cultures to PQ resulted in a significant ($p < 0.01$) and dose-dependent reduction in DA uptake (Fig. 1A), where 0.5 μ M and 1 μ M PQ caused 24% and 37% loss in DA neuronal function, respectively. To investigate the DA cell loss induced by PQ, neuron-glia cultures were exposed to PQ and stained with the TH antibody, and the number of TH-immunoreactive neurons was counted. PQ decreased the number of TH-immunoreactive neurons in a dose-dependent manner, with a 36% decrease in the cultures treated with 1 μ M PQ ($p < 0.01$) (Fig. 1B). In addition to the reduction in cell numbers, TH-immunoreactive neurons in PQ-treated cultures showed a less extensive dendritic network compared with those in control cultures (Fig. 1C).

PQ neurotoxicity is selective for DA neurons

To examine the selectivity of PQ neurotoxicity, [3 H]GABA uptake assay was performed to assess the function of GABAergic neurons in the cultures exposed to PQ. Figure 2 demonstrates that only DA uptake was reduced significantly ($p < 0.01$) by the addition of PQ to neuron-glia cultures, whereas GABA uptake remained unaffected ($p > 0.05$). LPS was used as a positive control for microglia activation and selective DA neurotoxicity (9).

Microglia mediate PQ-induced DA neurotoxicity

To investigate the role of microglia in PQ-induced DA neurotoxicity, [3 H]DA uptake was compared in microglia-depleted cultures and neuron-glia cultures. Higher doses of PQ (4 μ M and above) showed neurotoxicity in microglia-depleted cultures (data not shown), consistent with previous studies suggesting direct DA toxicity (39). However, whereas DA uptake was decreased in the neuron-glia culture exposed to 1 μ M PQ, there was no toxicity in microglia-depleted cultures ($p < 0.05$) (Fig. 3), suggesting the critical role of microglia in low-dose PQ neurotoxicity. LPS was used as a positive control for microglia-mediated DA neurotoxicity (9).

PQ induces extracellular $O_2^{\cdot-}$ production by microglia

It has been well documented that the activation of microglia and the consequent production of neurotoxic factors, including TNF α , nitric oxide, and ROS, have been linked to DA neurotoxicity (33, 37, 38). In an effort to discern the mechanism through which microglia exposed to PQ exerted neurotoxicity, the supernatant from PQ-treated neuron-glia cultures was tested for the presence of classic proinflammatory factors. Analysis of supernatant collected at 3 h, 6 h, 12 h, 24 h, 4 days, and 7 days post PQ treatment in neuron-glia cultures revealed that no TNF α or nitrite (indicative of nitric oxide production) was produced (data not shown). However, PQ did induce a dose-dependent increase in extracellular




FIG. 1. PQ is toxic to DA neurons. Mesencephalic midbrain neuron-glia cultures were treated with either vehicle, 0.5 μM PQ, or 1 μM PQ. (A) DA neurotoxicity was measured at 6–7 days post treatment using the [^3H]DA uptake assay. (B) DA cell death was determined at 6–7 days post treatment with immunocytochemical staining by using the TH antibody and counting the number of TH-positive neurons present. The data are expressed as the percentage of the control cultures and are the means \pm SEM from three independent experiments. An asterisk indicates a significant difference ($p < 0.01$) compared with control. (C) Cell morphology of TH-immunoreactive neurons in the neuron-glia cultures. The images shown are representative of three independent experiments.




FIG. 2. PQ neurotoxicity is selective for DA neurons. Mid-brain neuron-glia cultures were treated with either vehicle, 5 ng/ml LPS, or 1 μM PQ. LPS was used as a positive control for selective DA neurotoxicity. (A) DA neurotoxicity and GABA neurotoxicity were measured at 6–7 days post treatment by using the [^3H]DA uptake and [^3H]GABA uptake assays, respectively. The data are expressed as the percentage of the control cultures and are the means \pm SEM from three independent experiments. An asterisk indicates a significant difference ($p < 0.01$) compared with control.




FIG. 3. Microglia mediate DA neurotoxicity induced by 1 μM PQ. Mesencephalic neuron-glia cultures and microglia-depleted cultures were treated with either vehicle, 5 ng/ml LPS, or 1 μM PQ. LPS was used as a positive control for microglia-mediated DA neurotoxicity. DA neurotoxicity was measured at 6–7 days post treatment by using the [^3H]DA uptake assay. The data are expressed as the percentage of the control cultures and are the means \pm SEM from five independent experiments. An asterisk indicates a significant difference ($p < 0.05$) compared with control.




FIG. 4. PQ induces extracellular $O_2^{\cdot-}$ production in microglia. Primary microglia-enriched cultures were treated with either vehicle, 100 ng/ml LPS, 0.5 μ M PQ, or 1 μ M PQ. LPS was used as a positive control for microglial $O_2^{\cdot-}$ production. The production of extracellular $O_2^{\cdot-}$ was measured by the SOD-inhibitable reduction of tetrazolium salt, WST-1. The data are expressed as the percentage of the control cultures and are the means \pm SEM from five independent experiments. An asterisk indicates a significant difference ($p < 0.05$) compared with control.

$O_2^{\cdot-}$ production in enriched microglia ($p < 0.05$) (Fig. 4), which suggests that extracellular $O_2^{\cdot-}$ may be the main mediator of the microglia-mediated neurotoxicity. Consistent with previous reports (9, 33), LPS, as a positive control, was shown to induce $O_2^{\cdot-}$ production by microglia.

PQ-induced microglia-dependent neurotoxicity is mediated by NADPH oxidase

NADPH oxidase has been documented as the predominant source of extracellular $O_2^{\cdot-}$ in phagocytic cells. $PHOX^{-/-}$ mice lack the functional gp91 protein, the catalytic subunit of the NADPH oxidase complex, and thus produce no extracellular $O_2^{\cdot-}$ in response to immunological stimuli. DA uptake was compared between mesencephalic neuron-glia cultures from $PHOX^{-/-}$ mice and $PHOX^{+/+}$ mice. Figure 5 shows that whereas DA neurons in $PHOX^{+/+}$ cultures lost 36% of uptake function after 7-day treatment with 1 μ M PQ ($p < 0.01$), DA neurons in $PHOX^{-/-}$ cultures were completely resistant to the same dose of PQ. This suggests that, at 1 μ M, PQ neurotoxicity is mediated through NADPH oxidase and supports the importance of $O_2^{\cdot-}$ in DA neurotoxicity. LPS was used as a positive control for NADPH oxidase-mediated neurotoxicity, where LPS showed less DA neurotoxicity in midbrain neuron-glia cultures from $PHOX^{-/-}$ mice compared with $PHOX^{+/+}$ mice (33).

DISCUSSION

Oxidative stress has been strongly linked to neurodegenerative disease, with particular emphasis on PD. Thus, the identification of multiple, potential triggers of oxidative insult in




FIG. 5. NADPH oxidase mediates PQ DA neurotoxicity. $PHOX^{-/-}$ mice lack the functional catalytic subunit of the NADPH oxidase complex, gp91, and fail to produce the phagocytic respiratory burst. Mesencephalic neuron-glia cultures from $PHOX^{-/-}$ and $PHOX^{+/+}$ mice were treated with either vehicle, 5 ng/ml LPS, or 1 μ M PQ. LPS was used as a positive control for microglial NADPH oxidase activation. DA neurotoxicity was measured at 6–7 days post treatment using the [3 H]DA uptake assay. The data are expressed as the percentage of the control cultures and are the means \pm SEM from four independent experiments. An asterisk indicates a significant difference ($p < 0.01$) compared with control.

the brain and the elucidation of corresponding mechanisms are of paramount importance to both the understanding of disease progression and the development of neuroprotective compounds. In the present study, the neurotoxicity of PQ was examined in mesencephalic neuron-glia cultures, where 0.5–1 μ M PQ was shown to damage DA neurons in a dose-dependent manner (Fig. 1) and to be selectively toxic to DA neurons (Fig. 2). This is consistent with several previously reported *in vivo* studies (25, 26, 39), supporting the notion that PQ is indeed selectively toxic to DA neurons.

More interestingly, microglia were identified as the cell type governing PQ toxicity at lower concentrations (0.5–1 μ M). Originally, PQ was believed to be directly neurotoxic to DA neurons due to the structural similarity to MPP⁺ and because both are actively taken into the DA neuron through the DA transporter receptor (37, 38). Further, PQ is known to be able to redox-cycle with various diaphorases (3–5, 7) and oxygen to produce $O_2^{\cdot-}$. Thus, once inside the DA neurons, PQ may induce an increase of intracellular ROS and consequently damage the neurons. However, this study reports that whereas higher concentrations of PQ may indeed be directly toxic to DA neurons, lower concentrations (up to 1 μ M) require the presence of microglia in order to be toxic (Fig. 3). Microglia, the resident immune cells of the central nervous system, have been shown to play an important role in LPS- and rotenone-induced DA neuronal death, where microglial

NADPH oxidase-derived $O_2^{\cdot-}$ is the common mediator of cell death (8, 10, 33).

Similar to rotenone (40, 41), here we show that PQ activated microglia to produce ROS. The addition of PQ at any dose to the midbrain neuron-glia cultures failed to result in the production of soluble proinflammatory factors, a response commonly reported in the traditional microglial proinflammatory response. However, PQ did induce the production of extracellular $O_2^{\cdot-}$ from microglia (Fig. 4), identifying microglia as an additional source of PQ-induced oxidative stress. Although the exact species of ROS governing the neurotoxicity is unknown, there is a well established association between the microglial production of $O_2^{\cdot-}$ and DA neurotoxicity (35, 40, 41).

DA neurons are known to be particularly vulnerable to oxidative insult due to a reduced antioxidant capacity, a high content of iron and oxidation-prone DA, and a potential defect in mitochondrial complex I (12, 13). Thus, any compound inducing oxidative stress in the central nervous system is a potential candidate for selective DA neurotoxicity. However, the existence of a class of agrochemicals (rotenone and PQ) shown to induce oxidative stress at low concentrations solely through the production of microglial $O_2^{\cdot-}$ may provide novel insight into the mechanism governing the selective degeneration of DA neurons by brain region. Specifically, the SN is reported to contain 4.5 times as many microglia, compared with other regions of the brain (14), supporting the notion that the DA neurons localized there could be preferentially susceptible to microglia-mediated oxidative stress.

NADPH oxidase is the enzyme responsible for the microglial production of extracellular $O_2^{\cdot-}$ in response to LPS (35) and rotenone (40, 41). PHOX^{-/-} mice are devoid of the gp91 protein, catalytic subunit of NADPH oxidase complex, and are unable to produce extracellular $O_2^{\cdot-}$ in response to stimuli. In this study, we show that neuron-glia cultures from PHOX^{-/-} mice were also insensitive to 1 μ M PQ (Fig. 5), suggesting that NADPH oxidase is critical for PQ-induced neurotoxicity at lower concentrations. Current studies in our laboratory are focused on identifying the mechanisms of how PQ and rotenone activate NADPH oxidase in microglia.

In summary, we demonstrate that the selective DA neurotoxicity of lower concentrations (up to 1 μ M) of PQ is still mediated through the production of the $O_2^{\cdot-}$ radical. However, we report that at lower concentrations, the source of PQ-induced DA neurotoxic oxidative stress is initially extracellular and originates from the microglia, rather than from the neuron itself. Further, this microglia-mediated neurotoxicity is derived from the activation of NADPH oxidase. Together, these data suggest an emerging class of environmental agents (PQ and rotenone) that act at lower concentrations to provide an indirect source of oxidative insult through microglia, which is consequently selectively toxic to DA neurons.

ACKNOWLEDGMENTS

We thank Yu-Xiang Diao for his technical assistance. DISCLAIMER: This document has been reviewed by the National Health and Environmental Effects Research Laboratory and approved for publication. Approval does not signify

that the contents reflect the views of the Agency, nor does mention of trade names or commercial products constitute the endorsement of recommendation for use.

ABBREVIATIONS

DA, dopamine or dopaminergic; ELISA, enzyme-linked immunosorbent assay; GABA, γ -amino-*n*-butyric acid; HBSS, Hanks' balanced salt solution; LPS, lipopolysaccharide; MPP⁺, 1-methyl-4-phenylpyridinium; $O_2^{\cdot-}$, superoxide; PD, Parkinson's disease; PQ, paraquat; ROS, reactive oxygen species; SN, substantia nigra; SOD, superoxide dismutase; TH, tyrosine hydroxylase; TNF α , tumor necrosis factor- α .

REFERENCES

1. Barlow BK, Thiruchelvam MJ, Bennice L, Cory-Slechta DA, Ballatori N, and Richfield EK. Increased synaptosomal dopamine content and brain concentration of paraquat produced by selective dithiocarbamates. *J Neurochem* 85: 1075–1086, 2003.
2. Brooks AI, Chadwick CA, Gelbard HA, Cory-Slechta DA, and Federoff HJ. Paraquat elicited neurobehavioral syndrome caused by dopaminergic neuron loss. *Brain Res* 823: 1–10, 1999.
3. Bus JS, Aust SD, and Gibson JE. Superoxide- and singlet oxygen-catalyzed lipid peroxidation as a possible mechanism for paraquat (methyl viologen) toxicity. *Biochem Biophys Res Commun* 58: 749–755, 1974.
4. Clejan L and Cederbaum AI. Synergistic interactions between NADPH-cytochrome P-450 reductase, paraquat, and iron in the generation of active oxygen radicals. *Biochem Pharmacol* 38: 1779–1786, 1989.
5. Day BJ, Patel M, Calavetta L, Chang LY, and Stamler JS. A mechanism of paraquat toxicity involving nitric oxide synthase. *Proc Natl Acad Sci U S A* 96: 12760–12765, 1999.
6. Fredriksson A, Fredriksson M, and Eriksson P. Neonatal exposure to paraquat or MPTP induces permanent changes in striatum dopamine and behavior in adult mice. *Toxicol Appl Pharmacol* 122: 258–264, 1993.
7. Fukushima T, Yamada K, Isobe A, Shiwaku K, and Yamane Y. Mechanism of cytotoxicity of paraquat. I. NADH oxidation and paraquat radical formation via complex I. *Exp Toxicol Pathol* 45: 345–349, 1993.
8. Gao HM, Hong JS, Zhang W, and Liu B. Distinct role for microglia in rotenone-induced degeneration of dopaminergic neurons. *J Neurosci* 22: 782–790, 2002.
9. Gao HM, Jiang J, Wilson B, Zhang W, Hong JS, and Liu B. Microglial activation-mediated delayed and progressive degeneration of rat nigral dopaminergic neurons: relevance to Parkinson's disease. *J Neurochem* 81: 1285–1297, 2002.
10. Gao HM, Liu B, and Hong JS. Critical role for microglial NADPH oxidase in rotenone-induced degeneration of dopaminergic neurons. *J Neurosci* 23: 6181–6187, 2003.
11. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, and Tannenbaum SR. Analysis of nitrate, nitrite, and

- [¹⁵N]nitrate in biological fluids. *Anal Biochem* 126: 131–138, 1982.
12. Greenamyre JT, MacKenzie G, Peng TI, and Stephens SE. Mitochondrial dysfunction in Parkinson's disease. *Biochem Soc Symp* 66: 85–97, 1999.
 13. Jenner P and Olanow CW. Understanding cell death in Parkinson's disease. *Ann Neurol* 44: S72–S84, 1998.
 14. Kim WG, Mohny RP, Wilson B, Jeohn GH, Liu B, and Hong JS. Regional difference in susceptibility to lipopolysaccharide-induced neurotoxicity in the rat brain: role of microglia. *J Neurosci* 20: 6309–6316, 2000.
 15. Langston JW. Parkinson's disease: current and future challenges. *Neurotoxicology* 23: 443–450, 2002.
 16. Langston JW and Ballard PA Jr. Parkinson's disease in a chemist working with 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine. *N Engl J Med* 309: 310, 1983.
 17. Liou HH, Chen RC, Tsai YF, Chen WP, Chang YC, and Tsai MC. Effects of paraquat on the substantia nigra of the Wistar rats: neurochemical, histological, and behavioral studies. *Toxicol Appl Pharmacol* 137: 34–41, 1996.
 18. Liou HH, Tsai MC, Chen CJ, Jeng JS, Chang YC, Chen SY, and Chen RC. Environmental risk factors and Parkinson's disease: a case-control study in Taiwan. *Neurology* 48: 1583–1588, 1997.
 19. Liu B and Hong JS. Primary rat mesencephalic neuron-glia, neuron-enriched, microglia-enriched, and astroglia-enriched cultures. *Methods Mol Med* 79: 387–395, 2003.
 20. Liu B and Hong JS. Role of microglia in inflammation-mediated neurodegenerative diseases: mechanisms and strategies for therapeutic intervention. *J Pharmacol Exp Ther* 304: 1–7, 2003.
 21. Liu B, Du L, and Hong JS. Naloxone protects rat dopaminergic neurons against inflammatory damage through inhibition of microglia activation and superoxide generation. *J Pharmacol Exp Ther* 293: 607–617, 2000.
 22. Liu B, Qin L, Yang SN, Wilson BC, Liu Y, and Hong JS. Femtomolar concentrations of dynorphins protect rat mesencephalic dopaminergic neurons against inflammatory damage. *J Pharmacol Exp Ther* 298: 1133–1141, 2001.
 23. Liu B, Wang K, Gao HM, Mandavilli B, Wang JY, and Hong JS. Molecular consequences of activated microglia in the brain: overactivation induces apoptosis. *J Neurochem* 77: 182–189, 2001.
 24. Liu B, Gao HM, and Hong JS. Parkinson's disease and exposure to infectious agents and pesticides and the occurrence of brain injuries: role of neuroinflammation. *Environ Health Perspect* 111: 1065–1073, 2003.
 25. Manning-Bog AB, McCormack AL, Li J, Uversky VN, Fink AL, and Di Monte DA. The herbicide paraquat causes up-regulation and aggregation of alpha-synuclein in mice: paraquat and alpha-synuclein. *J Biol Chem* 277: 1641–1644, 2002.
 26. McCormack AL, Thiruchelvam M, Manning-Bog AB, Thiffault C, Langston JW, Cory-Slechta DA, and Di Monte DA. Environmental risk factors and Parkinson's disease: selective degeneration of nigral dopaminergic neurons caused by the herbicide paraquat. *Neurobiol Dis* 10: 119–127, 2002.
 27. McGeer PL and McGeer EG. Glial cell reactions in neurodegenerative diseases: pathophysiology and therapeutic interventions. *Alzheimer Dis Assoc Disord* 12 Suppl 2: S1–S6, 1998.
 28. Olanow CW and Tatton WG. Etiology and pathogenesis of Parkinson's disease. *Annu Rev Neurosci* 22: 123–144, 1999.
 29. Perry TL, Yong VW, Wall RA, and Jones K. Paraquat and two endogenous analogues of the neurotoxic substance *N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine do not damage dopaminergic nigrostriatal neurons in the mouse. *Neurosci Lett* 69: 285–289, 1986.
 30. Peskin AV and Winterbourn CC. A microtiter plate assay for superoxide dismutase using a water-soluble tetrazolium salt (WST-1). *Clin Chim Acta* 293: 157–166, 2000.
 31. Priyadarshi A, Khuder SA, Schaub EA, and Priyadarshi SS. Environmental risk factors and Parkinson's disease: a metaanalysis. *Environ Res* 86: 122–127, 2001.
 32. Qin L, Liu Y, Cooper C, Liu B, Wilson B, and Hong JS. Microglia enhance beta-amyloid peptide-induced toxicity in cortical and mesencephalic neurons by producing reactive oxygen species. *J Neurochem* 83: 973–983, 2002.
 33. Qin L, Liu Y, Wang T, Wei SJ, Block ML, Wilson B, Liu B, and Hong JS. NADPH oxidase mediates lipopolysaccharide-induced neurotoxicity and proinflammatory gene expression in activated microglia. *J Biol Chem* 279: 1415–1421, 2004.
 34. Rajput AH and Uitti RJ. Paraquat and Parkinson's disease. *Neurology* 37: 1820–1821, 1987.
 35. Rajput AH, Uitti RJ, Stern W, Laverty W, O'Donnell K, O'Donnell D, Yuen WK, and Dua A. Geography, drinking water chemistry, pesticides and herbicides and the etiology of Parkinson's disease. *Can J Neurol Sci* 14: 414–418, 1987.
 36. Semchuk KM, Love EJ, and Lee RG. Parkinson's disease and exposure to agricultural work and pesticide chemicals. *Neurology* 42: 1328–1335, 1992.
 37. Shimizu K, Ohtaki K, Matsubara K, Aoyama K, Uezono T, Saito O, Suno M, Ogawa K, Hayase N, Kimura K, and Shiono H. Carrier-mediated processes in blood-brain barrier penetration and neural uptake of paraquat. *Brain Res* 906: 135–142, 2001.
 38. Shimizu K, Matsubara K, Ohtaki K, Fujimaru S, Saito O, and Shiono H. Paraquat induces long-lasting dopamine overflow through the excitotoxic pathway in the striatum of freely moving rats. *Brain Res* 976: 243–252, 2003.
 39. Shimizu K, Matsubara K, Ohtaki K, and Shiono H. Paraquat leads to dopaminergic neural vulnerability in organotypic midbrain culture. *Neurosci Res* 46: 523–532, 2003.
 40. Tan AS and Berridge MV. Superoxide produced by activated neutrophils efficiently reduces the tetrazolium salt, WST-1 to produce a soluble formazan: a simple colorimetric assay for measuring respiratory burst activation and for screening anti-inflammatory agents. *J Immunol Methods* 238: 59–68, 2000.
 41. Tawara T, Fukushima T, Hojo N, Isobe A, Shiwaku K, Setogawa T, and Yamane Y. Effects of paraquat on mitochondrial electron transport system and catecholamine contents in rat brain. *Arch Toxicol* 70: 585–589, 1996.
 42. Teismann P, Tieu K, Choi DK, Wu DC, Naini A, Hunot S, Vila M, Jackson-Lewis V, and Przedborski S. Cyclooxygenase-2 is instrumental in Parkinson's disease neurodegeneration. *Proc Natl Acad Sci USA* 100: 5473–5478, 2003.

43. Thiruchelvam M, Richfield EK, Baggs RB, Tank AW, and Cory-Slechta DA. The nigrostriatal dopaminergic system as a preferential target of repeated exposures to combined paraquat and maneb: implications for Parkinson's disease. *J Neurosci* 20: 9207–9214, 2000.
44. Uversky VN, Li J, and Fink AL. Pesticides directly accelerate the rate of alpha-synuclein fibril formation: a possible factor in Parkinson's disease. *FEBS Lett* 500: 105–108, 2001.
45. Widdowson PS, Farnworth MJ, Upton R, and Simpson MG. No changes in behaviour, nigro-striatal system neurochemistry or neuronal cell death following toxic multiple oral paraquat administration to rats. *Hum Exp Toxicol* 15: 583–591, 1996.
46. Woolley DE, Gietzen DW, Gee SJ, Magdalou J, and Hammock BD. Does paraquat (PQ) mimic MPP⁺ toxicity? *Proc West Pharmacol Soc* 32: 191–193, 1989.
47. Yumino K, Kawakami I, Tamura M, Hayashi T, and Nakamura M. Paraquat- and diquat-induced oxygen radical generation and lipid peroxidation in rat brain microsomes. *J Biochem (Tokyo)* 131: 565–570, 2002.

Address reprint requests to:
Jau-Shyong Hong, Ph.D.
MD F1-01 NIEHS
P.O. Box 12233
Research Triangle Park, NC 27709

E-mail: hong3@niehs.nih.gov

Received for publication March 24, 2004; accepted November 30, 2004.

This article has been cited by:

1. Maria Teresa Baltazar, Ricardo Jorge Dinis-Oliveira, Maria de Lourdes Bastos, Aristidis M. Tsatsakis, José Alberto Duarte, Félix Carvalho. 2014. Pesticides exposure as etioloical factors of Parkinson's disease and other neurodegenerative diseases—A mechanistic approach. *Toxicology Letters* . [[CrossRef](#)]
2. Jenny Sandström von Tobel, Debora Zoia, Jorane Althaus, Paola Antinori, Julien Mermoud, Hui Song Pak, Alexander Scherl, Florianne Monnet-Tschudi. 2014. Immediate and delayed effects of subchronic Paraquat exposure during an early differentiation stage in 3D-rat brain cell cultures. *Toxicology Letters* . [[CrossRef](#)]
3. Emanuele D'Amico, Pam Factor-Litvak, Regina M. Santella, Hiroshi Mitsumoto. 2013. Clinical perspective on oxidative stress in sporadic amyotrophic lateral sclerosis. *Free Radical Biology and Medicine* **65**, 509-527. [[CrossRef](#)]
4. Juliana Navarro-Yepes, Laura Zavala-Flores, Anandhan Annadurai, Fang Wang, Maciej Skotak, Namas Chandra, Ming Li, Aglaia Pappa, Daniel Martinez-Fong, Luz Maria Del Razo, Betzabet Quintanilla-Vega, Rodrigo Franco. 2013. Antioxidant gene therapy against neuronal cell death. *Pharmacology & Therapeutics* . [[CrossRef](#)]
5. Hjalte Holm Andersen, Kasper Bendix Johnsen, Torben Moos. 2013. Iron deposits in the chronically inflamed central nervous system and contributes to neurodegeneration. *Cellular and Molecular Life Sciences* . [[CrossRef](#)]
6. Xiuli Chang, Wen Lu, Tingting Dou, Xinjin Wang, Dan Lou, Xia Sun, Zhijun Zhou. 2013. Paraquat inhibits cell viability via enhanced oxidative stress and apoptosis in human neural progenitor cells. *Chemico-Biological Interactions* **206**:2, 248-255. [[CrossRef](#)]
7. Claire J. Williams, David T. Dexter. 2013. Neuroprotective and symptomatic effects of targeting group III mGlu receptors in neurodegenerative disease. *Journal of Neurochemistry* n/a-n/a. [[CrossRef](#)]
8. Humberto Rodriguez-Rocha, Aracely Garcia-Garcia, Chillian Pickett, Sumin Li, Jocelyn Jones, Han Chen, Brian Webb, Jae Choi, You Zhou, Matthew C. Zimmerman, Rodrigo Franco. 2013. Compartmentalized oxidative stress in dopaminergic cell death induced by pesticides and complex I inhibitors: Distinct roles of superoxide anion and superoxide dismutases. *Free Radical Biology and Medicine* **61**, 370-383. [[CrossRef](#)]
9. Manjeet Singh, Ven Murthy, Charles Ramassamy. 2013. Neuroprotective mechanisms of the standardized extract of Bacopa monniera in a paraquat/diquat-mediated acute toxicity. *Neurochemistry International* **62**:5, 530-539. [[CrossRef](#)]
10. Bailin Wu, Bo Song, Haiqing Yang, Boyuan Huang, Bo Chi, Yansu Guo, Huaijun Liu. 2013. Central nervous system damage due to acute paraquat poisoning: An experimental study with rat model. *NeuroToxicology* **35**, 62-70. [[CrossRef](#)]
11. Thomas Taetzsch, Michelle L. Block. 2013. Pesticides, Microglial NOX2, and Parkinson's Disease. *Journal of Biochemical and Molecular Toxicology* **27**:2, 137-149. [[CrossRef](#)]
12. Oddvar Myhre, Hans Utkilen, Nur Duale, Gunnar Brunborg, Tim Hofer. 2013. Metal Dyshomeostasis and Inflammation in Alzheimer's and Parkinson's Diseases: Possible Impact of Environmental Exposures. *Oxidative Medicine and Cellular Longevity* **2013**, 1-19. [[CrossRef](#)]
13. Sandeep Vasant More, Hemant Kumar, In Su Kim, Soo-Yeol Song, Dong-Kug Choi. 2013. Cellular and Molecular Mediators of Neuroinflammation in the Pathogenesis of Parkinson's Disease. *Mediators of Inflammation* **2013**, 1-12. [[CrossRef](#)]
14. Aracely Garcia-Garcia, Laura Zavala-Flores, Humberto Rodriguez-Rocha, Rodrigo Franco. 2012. Thiol-Redox Signaling, Dopaminergic Cell Death, and Parkinson's Disease. *Antioxidants & Redox Signaling* **17**:12, 1764-1784. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
15. Ailing Ji, Huan Diao, Xiaohui Wang, Ruihua Yang, Jianbin Zhang, Wenjing Luo, Rui Cao, Zipeng Cao, Feng Wang, Tongjian Cai. 2012. n-3 polyunsaturated fatty acids inhibit lipopolysaccharide-induced microglial activation and dopaminergic injury in rats. *NeuroToxicology* **33**:4, 780-788. [[CrossRef](#)]
16. J. Bové, C. Perier. 2012. Neurotoxin-based models of Parkinson's disease. *Neuroscience* **211**, 51-76. [[CrossRef](#)]
17. Hui-Ming Gao, Hui Zhou, Jau-Shyong Hong. 2012. NADPH oxidases: novel therapeutic targets for neurodegenerative diseases. *Trends in Pharmacological Sciences* **33**:6, 295-303. [[CrossRef](#)]
18. Rommy von Bernhardt, Jaime Eugenin. 2012. Alzheimer's Disease: Redox Dysregulation As a Common Denominator for Diverse Pathogenic Mechanisms. *Antioxidants & Redox Signaling* **16**:9, 974-1031. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
19. Xiao-Guang Luo, Sheng-Di Chen. 2012. The changing phenotype of microglia from homeostasis to disease. *Translational Neurodegeneration* **1**:1, 9. [[CrossRef](#)]

20. Sushruta Koppula, Hemant Kumar, In Su Kim, Dong-Kug Choi. 2012. Reactive Oxygen Species and Inhibitors of Inflammatory Enzymes, NADPH Oxidase, and iNOS in Experimental Models of Parkinson's Disease. *Mediators of Inflammation* **2012**, 1-16. [[CrossRef](#)]
21. Sharawan Yadav, Satya Prakash Gupta, Garima Srivastava, Pramod Kumar Srivastava, Mahendra Pratap Singh. 2011. Role of Secondary Mediators in Caffeine-Mediated Neuroprotection in Maneb- and Paraquat-Induced Parkinson's Disease Phenotype in the Mouse. *Neurochemical Research* . [[CrossRef](#)]
22. P. M. Rappold, M. Cui, A. S. Chesser, J. Tibbett, J. C. Grima, L. Duan, N. Sen, J. A. Javitch, K. Tieu. 2011. Paraquat neurotoxicity is mediated by the dopamine transporter and organic cation transporter-3. *Proceedings of the National Academy of Sciences* **108**:51, 20766-20771. [[CrossRef](#)]
23. Ashutosh Kumar, Brajesh Kumar Singh, Israr Ahmad, Smriti Shukla, Devendra Kumar Patel, Garima Srivastava, Vinod Kumar, Haushila Prasad Pandey, Chetna Singh. 2011. Involvement of NADPH oxidase and glutathione in zinc-induced dopaminergic neurodegeneration in rats: Similarity with paraquat neurotoxicity. *Brain Research* . [[CrossRef](#)]
24. M. Singh, V. Murthy, C. Ramassamy. 2011. Standardized extracts of Bacopa monniera protect against MPP+ and Paraquat induced- toxicities by modulating mitochondrial activities, proteasomal functions and redox pathways. *Toxicological Sciences* . [[CrossRef](#)]
25. Che Brown Hutson, Carlos R. Lazo, Farzad Mortazavi, Christopher C. Giza, David Hovda, Marie-Francoise Chesselet. 2011. Traumatic Brain Injury in Adult Rats Causes Progressive Nigrostriatal Dopaminergic Cell Loss and Enhanced Vulnerability to the Pesticide Paraquat. *Journal of Neurotrauma* **28**:9, 1783-1801. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
26. Hui-Ming Gao, Jau-Shyong Hong. 2011. Gene-environment interactions: Key to unraveling the mystery of Parkinson's disease. *Progress in Neurobiology* **94**:1, 1-19. [[CrossRef](#)]
27. Darcy Litteljohn, Emily Mangano, Melanie Clarke, Jessica Boby, Kerry Moloney, Shawn Hayley. 2011. Inflammatory Mechanisms of Neurodegeneration in Toxin-Based Models of Parkinson's Disease. *Parkinson's Disease* **2011**, 1-18. [[CrossRef](#)]
28. Rodrigo Franco, Sumin Li, Humberto Rodriguez-Rocha, Michaela Burns, Mihalis I. Panayiotidis. 2010. Molecular mechanisms of pesticide-induced neurotoxicity: Relevance to Parkinson's disease. *Chemico-Biological Interactions* **188**:2, 289-300. [[CrossRef](#)]
29. Elisabetta Polazzi, Barbara Monti. 2010. Microglia and neuroprotection: From in vitro studies to therapeutic applications. *Progress in Neurobiology* **92**:3, 293-315. [[CrossRef](#)]
30. Melinda E. Lull, Michelle L. Block. 2010. Microglial Activation and Chronic Neurodegeneration. *Neurotherapeutics* **7**:4, 354-365. [[CrossRef](#)]
31. M. A. Ortiz-Ortiz, J. M. Moran, L. M. Ruiz-Mesa, J. M. Bravo-San Pedro, J. M. Fuentes. 2010. Paraquat Exposure Induces Nuclear Translocation of Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH) and the Activation of the Nitric Oxide-GAPDH-Siah Cell Death Cascade. *Toxicological Sciences* **116**:2, 614-622. [[CrossRef](#)]
32. Sergio Rosales-Corral, Russel J. Reiter, Dun-Xian Tan, Genaro G. Ortiz, Gabriela Lopez-Armas. 2010. Functional Aspects of Redox Control During Neuroinflammation. *Antioxidants & Redox Signaling* **13**:2, 193-247. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
33. C Berry, C La Vecchia, P Nicotera. 2010. Paraquat and Parkinson's disease. *Cell Death and Differentiation* **17**:7, 1115-1125. [[CrossRef](#)]
34. Christopher J. Barnum, Malú G. Tansey Modeling neuroinflammatory pathogenesis of Parkinson's disease **184**, 113-132. [[CrossRef](#)]
35. Woody McGinnis, Veronica Miller, Tapan Audhya, Stephen Edelson Neurotoxic Brainstem Impairment as Proposed Threshold Event in Autistic Regression 153-176. [[CrossRef](#)]
36. Ana Clara Cristóvão, Dong-Hee Choi, Graça Baltazar, M. Flint Beal, Yoon-Seong Kim. 2009. The Role of NADPH Oxidase 1-Derived Reactive Oxygen Species in Paraquat-Mediated Dopaminergic Cell Death. *Antioxidants & Redox Signaling* **11**:9, 2105-2118. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
37. Kavita Prasad, Elizabeth Tarasewicz, Jason Mathew, Pamela A. Ohman Strickland, Brian Buckley, Jason R. Richardson, Eric K. Richfield. 2009. Toxicokinetics and toxicodynamics of paraquat accumulation in mouse brain. *Experimental Neurology* **215**:2, 358-367. [[CrossRef](#)]
38. Rebecca L. Miller, Marilyn James-Kracke, Grace Y. Sun, Albert Y. Sun. 2009. Oxidative and Inflammatory Pathways in Parkinson's Disease. *Neurochemical Research* **34**:1, 55-65. [[CrossRef](#)]

39. Jun Peng, Fang Feng Stevenson, May Lin Oo, Julie K. Andersen. 2009. Iron-enhanced paraquat-mediated dopaminergic cell death due to increased oxidative stress as a consequence of microglial activation. *Free Radical Biology and Medicine* **46**:2, 312-320. [[CrossRef](#)]
40. Albert Y. Sun, Qun Wang, Agnes Simonyi, Grace Y. Sun. 2008. Botanical Phenolics and Brain Health. *NeuroMolecular Medicine* **10**:4, 259-274. [[CrossRef](#)]
41. Hui-Ming Gao, Jau-Shyong Hong. 2008. Why neurodegenerative diseases are progressive: uncontrolled inflammation drives disease progression. *Trends in Immunology* **29**:8, 357-365. [[CrossRef](#)]
42. Sangseop Kim, Jaegyung Hwang, Won-Ha Lee, Dae Youn Hwang, Kyoungsook Suk. 2008. Role of protein kinase C δ in paraquat-induced glial cell death. *Journal of Neuroscience Research* **86**:9, 2062-2070. [[CrossRef](#)]
43. Derek A. Drechsel, Manisha Patel. 2008. Role of reactive oxygen species in the neurotoxicity of environmental agents implicated in Parkinson's disease. *Free Radical Biology and Medicine* **44**:11, 1873-1886. [[CrossRef](#)]
44. Jung-Sung Chung, Jian-Kang Zhu, Ray A. Bressan, Paul M. Hasegawa, Huazhong Shi. 2008. Reactive oxygen species mediate Na⁺-induced SOS1 mRNA stability in Arabidopsis. *The Plant Journal* **53**:3, 554-565. [[CrossRef](#)]
45. Kathleen A. Maguire-Zeiss, Xiaomin Su, Howard J. Federoff. Microglial activation in a mouse model of α -synuclein overexpression 545-557. [[CrossRef](#)]
46. Rebecca L. Miller, Grace Y. Sun, Albert Y. Sun. 2007. Cytotoxicity of paraquat in microglial cells: Involvement of PKC δ - and ERK1/2-dependent NADPH oxidase. *Brain Research* **1167**, 129-139. [[CrossRef](#)]
47. Thomas C. Long, Julianne Tajuba, Preethi Sama, Navid Saleh, Carol Swartz, Joel Parker, Susan Hester, Gregory V. Lowry, Bellina Veronesi. 2007. Nanosize Titanium Dioxide Stimulates Reactive Oxygen Species in Brain Microglia and Damages Neurons in Vitro. *Environmental Health Perspectives* **115**:11, 1631-1637. [[CrossRef](#)]
48. P S Whitton. 2007. Inflammation as a causative factor in the aetiology of Parkinson's disease. *British Journal of Pharmacology* **150**:8, 963-976. [[CrossRef](#)]
49. Liya Qin, Xuefei Wu, Michelle L. Block, Yuxin Liu, George R. Breese, Jau-Shyong Hong, Darin J. Knapp, Fulton T. Crews. 2007. Systemic LPS causes chronic neuroinflammation and progressive neurodegeneration. *Glia* **55**:5, 453-462. [[CrossRef](#)]
50. Karl-Heinz Krause. 2007. Aging: A revisited theory based on free radicals generated by NOX family NADPH oxidases. *Experimental Gerontology* **42**:4, 256-262. [[CrossRef](#)]
51. Michelle L. Block, Luigi Zecca, Jau-Shyong Hong. 2007. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nature Reviews Neuroscience* **8**:1, 57-69. [[CrossRef](#)]
52. P-S Chen, G-S Peng, G Li, S Yang, X Wu, C-C Wang, B Wilson, R-B Lu, P-W Gean, D-M Chuang, J-S Hong. 2006. Valproate protects dopaminergic neurons in midbrain neuron/glia cultures by stimulating the release of neurotrophic factors from astrocytes. *Molecular Psychiatry* **11**:12, 1116-1125. [[CrossRef](#)]
53. Weihua Zhao, Wenjie Xie, Qin Xiao, David R. Beers, Stanley H. Appel. 2006. Protective effects of an anti-inflammatory cytokine, interleukin-4, on motoneuron toxicity induced by activated microglia. *Journal of Neurochemistry* **99**:4, 1176-1187. [[CrossRef](#)]
54. Bin Liu. 2006. Modulation of microglial pro-inflammatory and neurotoxic activity for the treatment of Parkinson's disease. *The AAPS Journal* **8**:3, E606-E621. [[CrossRef](#)]
55. Yulia Y. Tyurina, Alexander A. Kapralov, Jianfei Jiang, Grigory G. Borisenko, Alla I. Potapovich, Andrey Sorokin, Patrick M. Kochanek, Steven H. Graham, Nina F. Schor, Valerian E. Kagan. 2006. Oxidation and cytotoxicity of 6-OHDA are mediated by reactive intermediates of COX-2 overexpressed in PC12 cells. *Brain Research* **1093**:1, 71-82. [[CrossRef](#)]
56. Liya Qin, Guorong Li, Xun Qian, Yuxin Liu, Xuefei Wu, Bin Liu, Jau-Shyong Hong, Michelle L. Block. 2005. Interactive role of the toll-like receptor 4 and reactive oxygen species in LPS-induced microglia activation. *Glia* **52**:1, 78-84. [[CrossRef](#)]
57. LIYA QIN, YUXIN LIU, XUN QIAN, JAU-SHYONG HONG, MICHELLE L. BLOCK. 2005. Microglial NADPH Oxidase Mediates Leucine Enkephalin Dopaminergic Neuroprotection. *Annals of the New York Academy of Sciences* **1053**:1, 107-120. [[CrossRef](#)]
58. Michelle L. Block, Jau-Shyong Hong. 2005. Microglia and inflammation-mediated neurodegeneration: Multiple triggers with a common mechanism. *Progress in Neurobiology* **76**:2, 77-98. [[CrossRef](#)]
59. Todd B. Sherer, J. Timothy Greenamyre. 2005. Oxidative Damage in Parkinson's Disease. *Antioxidants & Redox Signaling* **7**:5-6, 627-629. [[Citation](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]